

# Intuitionistic Fuzzy Multi Similarity Measure Based on Cotangent Function

\* P. Rajarajeswari,    \*\* N. Uma

\* Department of Mathematics, Chikkanna Arts College, Tirupur, Tamil Nadu. (INDIA).

\*\* Department of Mathematics, SNR Sons College, Coimbatore, Tamil Nadu. (INDIA).

## ABSTRACT

*In this paper, the Similarity measure of Intuitionistic Fuzzy Multi sets (IFMS) based on Cotangent function is presented and analyzed. The properties of the similarity measure are proved and verified using the numerical evaluation of the proposed similarity measure. The unique feature of this proposed method is that it considers multi membership and non membership for the same element. As the proposed method is mathematically valid, it can be applied to any decision making problems, medical diagnosis, engineering problems, pattern recognition, etc. Finally, the application of medical diagnosis shows that the proposed similarity measures are much simpler, well suited one to use with linguistic variables.*

**KEY WORDS:** Intuitionistic fuzzy set, Intuitionistic Fuzzy Multi sets, Similarity measure, Cotangent Function

## INTRODUCTION

**Krassimir T. Atanassov** [1], [2] proposed the Intuitionistic Fuzzy sets (IFS) as the generalisation of the Fuzzy set (FS) introduced by **Lofti A. Zadeh** [3]. The FS allows the object to partially belong to a set with a membership degree ( $\mu$ ) between 0 and 1 whereas IFS represent the uncertainty with respect to both membership ( $\mu \in [0,1]$ ) and non membership ( $\vartheta \in [0,1]$ ) such that  $\mu + \vartheta \leq 1$ . The number  $\pi = 1 - \mu - \vartheta$  is called the hesitation degree or intuitionistic index.

The Multi set [4] allows the repeated occurrences of any element and hence the Fuzzy Multi set (FMS) can occur more than once with the possibly of the same or the different membership values was introduced by **R. R. Yager** [5]. Recently, the new concept Intuitionistic Fuzzy Multi sets (IFMS) was proposed by **T.K Shinoj and Sunil Jacob John** [6].

The study of distance and similarity measure of IFSs gives lots of measures, each representing specific properties and behaviour in real-life decision making and pattern recognition works. For measuring the degree of similarity between vague sets, **Chen and Tan** [7] proposed two similarity measures. The Hamming, Euclidean distance and similarity measures were introduced by **Szmidt and Kacprzyk** [8], [9], [10], [11]. The Geometric distance and similarity measures were given by **Xu** (12). Using the Cotangent function, a new similarity measure was proposed by **Wang et al** (13). Later a new fuzzy cotangent similarity measure for IFSs was introduced by **Tian Maoying** (14).

As the extension of the distance and similarity measure of IFSs to IFMSs [15], [16] are possible; in this paper we extend the fuzzy cotangent similarity measure of IFSs to IFMSs. The numerical results of the examples show that the developed similarity measures are well suited to use any linguistic variables.

The organization of this paper is as follows: In section 2, the Fuzzy Multi sets, Intuitionistic Fuzzy Multi sets and similarity measures of IFMS are presented. The section 3 deals with the proposed Cotangent Similarity measure for the IFMS, along with the numerical evaluation. The application of medical diagnosis using IFMS is explained in detail in section 4.

## II. PRELIMINARIES

### Definition: 2.1

Let  $X$  be a nonempty set. A **fuzzy set**  $A$  in  $X$  is given by  $A = \{(x, \mu_A(x)) / x \in X\}$  -- (2.1)

where  $\mu_A : X \rightarrow [0, 1]$  is the membership function of the fuzzy set  $A$  (i.e.)  $\mu_A(x) \in [0,1]$  is the membership of  $x \in X$  in  $A$ . The generalizations of fuzzy sets are the Intuitionistic fuzzy (IFS) set proposed by **Atanassov** [1], [2] is with independent memberships and non memberships.

**Definition: 2.2**

An *Intuitionistic fuzzy set (IFS)*,  $A$  in  $X$  is given by  $A = \{ \langle x, \mu_A(x), \vartheta_A(x) \rangle / x \in X \}$  -- (2.2)

where  $\mu_A : X \rightarrow [0,1]$  and  $\vartheta_A : X \rightarrow [0,1]$  with the condition  $0 \leq \mu_A(x) + \vartheta_A(x) \leq 1$ ,  $\forall x \in X$ . Here  $\mu_A(x)$  and  $\vartheta_A(x) \in [0,1]$  denote the membership and the non membership functions of the fuzzy set  $A$ ; For each Intuitionistic fuzzy set in  $X$ ,  $\pi_A(x) = 1 - \mu_A(x) - \vartheta_A(x) = 0$  for all  $x \in X$  that is

$\pi_A(x) = 1 - \mu_A(x) - \vartheta_A(x)$  is the hesitancy degree of  $x \in X$  in  $A$ . Always  $0 \leq \pi_A(x) \leq 1$ ,  $\forall x \in X$ .

The *complementary set*  $A^c$  of  $A$  is defined as  $A^c = \{ \langle x, \vartheta_A(x), \mu_A(x) \rangle / x \in X \}$  -- (2.3)

**Definition: 2.3**

Let  $X$  be a nonempty set. A *Fuzzy Multi set (FMS)*  $A$  in  $X$  is characterized by the count membership function  $Mc$  such that  $Mc : X \rightarrow Q$  where  $Q$  is the set of all crisp multi sets in  $[0,1]$ . Hence, for any  $x \in X$ ,  $Mc(x)$  is the crisp multi set from  $[0, 1]$ . The membership sequence is defined as

$(\mu_A^1(x), \mu_A^2(x), \dots, \mu_A^p(x))$  where  $\mu_A^1(x) \geq \mu_A^2(x) \geq \dots \geq \mu_A^p(x)$ .

Therefore, A FMS  $A$  is given by  $A = \{ \langle x, (\mu_A^1(x), \mu_A^2(x), \dots, \mu_A^p(x)) \rangle / x \in X \}$  -- (2.4)

**Definition: 2.4**

Let  $X$  be a nonempty set. A *Intuitionistic Fuzzy Multi set (IFMS)*  $A$  in  $X$  is characterized by two functions namely count membership function  $Mc$  and count non membership function  $NMc$  such that  $Mc : X \rightarrow Q$  and  $NMc : X \rightarrow Q$  where  $Q$  is the set of all crisp multi sets in  $[0,1]$ . Hence, for any  $x \in X$ ,  $Mc(x)$  is the crisp multi set from  $[0, 1]$  whose membership sequence is defined as

$(\mu_A^1(x), \mu_A^2(x), \dots, \mu_A^p(x))$  where  $\mu_A^1(x) \geq \mu_A^2(x) \geq \dots \geq \mu_A^p(x)$  and the corresponding

non membership sequence  $NMc(x)$  is defined as  $(\vartheta_A^1(x), \vartheta_A^2(x), \dots, \vartheta_A^p(x))$  where the non membership can be either decreasing or increasing function. such that  $0 \leq \mu_A^i(x) + \vartheta_A^i(x) \leq 1$ ,  $\forall x \in X$  and  $i = 1, 2, \dots, p$ . Therefore,

An *IFMS*  $A$  is given by  $A = \{ \langle x, (\mu_A^1(x), \mu_A^2(x), \dots, \mu_A^p(x)), (\vartheta_A^1(x), \vartheta_A^2(x), \dots, \vartheta_A^p(x)) \rangle / x \in X \}$  -- (2.5)

where  $\mu_A^1(x) \geq \mu_A^2(x) \geq \dots \geq \mu_A^p(x)$ . The *complementary set*  $A^c$  of  $A$  is defined as

$A^c = \{ \langle x, (\vartheta_A^1(x), \vartheta_A^2(x), \dots, \vartheta_A^p(x)), (\mu_A^1(x), \mu_A^2(x), \dots, \mu_A^p(x)) \rangle / x \in X \}$  -- (2.6)

where  $\vartheta_A^1(x) \geq \vartheta_A^2(x) \geq \dots \geq \vartheta_A^p(x)$

**Definition: 2.5**

The **Cardinality** of the membership function  $Mc(x)$  and the non membership function  $NMc(x)$  is the length of an element  $x$  in an *IFMS*  $A$  denoted as  $\eta$ , defined as  $\eta = |Mc(x)| = |NMc(x)|$

If  $A, B, C$  are the *IFMS* defined on  $X$ , then their cardinality  $\eta = \text{Max} \{ \eta(A), \eta(B), \eta(C) \}$ .

**Definition: 2.6**

*Sim* ( $A, B$ ) is said to be the **similarity measure** between  $A$  and  $B$ , where  $A, B \in X$  and  $X$  is an *IFMS*, as *Sim* ( $A, B$ ) satisfies the following properties

1.  $Sim(A, B) \in [0,1]$
2.  $Sim(A, B) = 1$  if and only if  $A = B$
3.  $Sim(A, B) = Sim(B, A)$
4. If  $A \subseteq B \subseteq C \in X$ , then  $Sim(A, C) \leq Sim(A, B)$  and  $Sim(A, C) \leq Sim(B, C)$

### COTANGENT SIMILARITY MEASURE OF *IFS*s

The entropy measure of *IFS*s proposed by Wang et al [13] was as follows

$$E(A) = \frac{1}{n} \sum_{i=1}^n \text{Cot} \left\{ \frac{1}{4} \pi + \frac{|\mu_A(x) - \vartheta_A(x)| \pi}{4(1 + \pi_A(x))} \right\}$$

As there is a relationship between the Entropy and Similarity Measure, the fuzzy cotangent similarity measure of *IFS*s was proposed by Tian Maoying [14]

$$FCS(A, B) = \frac{1}{n} \sum_{i=1}^n \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A(x_i) - \mu_B(x_i)| \vee |\vartheta_A(x_i) - \vartheta_B(x_i)|)}{4} \right\}$$

where  $A, B \in \text{IFS}(X)$  and  $E$  be the entropy of *IFS*s, consisting of the membership and non membership functions. If there are three parameters like membership, non membership and hesitation function then the fuzzy cotangent similarity measure of *IFS*s becomes

$$FCS(A, B) = \frac{1}{n} \sum_{i=1}^n \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A(x_i) - \mu_B(x_i)| \vee |\vartheta_A(x_i) - \vartheta_B(x_i)| \vee |\pi_A(x_i) - \pi_B(x_i)|)}{4} \right\}$$

### III PROPOSED COTANGENT SIMILARITY MEASURE OF *IFMS*s

In *IFS*, the Similarity measures are considered for the membership and non membership functions only once. But in *IFMS*, it should be considered more than once; because of their multi membership and non membership functions. And, their considerations are combined together by means of Summation concept based on their cardinality.

**Definition: 3.1**

$$IFMS(A, B) = \frac{1}{\eta} \sum_{j=1}^{\eta} \left[ \frac{1}{n} \sum_{i=1}^n \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A^j(x_i) - \mu_B^j(x_i)| \vee |\vartheta_A^j(x_i) - \vartheta_B^j(x_i)|)}{4} \right\} \right]$$

of the membership and non membership functions. And if there are three parameters like membership, non membership and hesitation function then the fuzzy multi cotangent similarity measure becomes

$$IFMS(A, B) = \frac{1}{\eta} \sum_{j=1}^{\eta} \left[ \frac{1}{n} \sum_{i=1}^n \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A^j(x_i) - \mu_B^j(x_i)| \vee |\vartheta_A^j(x_i) - \vartheta_B^j(x_i)| \vee |\pi_A^j(x_i) - \pi_B^j(x_i)|)}{4} \right\} \right]$$

### PROPOSITION: 3.2

The defined similarity measure  $IFMS(A, B)$  between *IFMS*  $A$  and  $B$  satisfies the following properties

- D1.  $0 \leq IFMS(A, B) \leq 1$
- D2.  $A = B$  if and only if  $IFMS(A, B) = 1$
- D3.  $IFMS(A, B) = IFMS(B, A)$
- D4. If  $A \subseteq B \subseteq C$ , for  $A, B, C$  are *IFMS* then

$$IFMS(A, B) \leq IFMS(A, C) \text{ and } IFMS(B, C) \leq IFMS(A, C)$$

**Proof****D1.  $0 \leq IFMS(A, B) \leq 1$** 

As the membership and the non membership functions of the *IFMSs* lies between 0 and 1, the similarity measure based cotangent function also lies between 0 and 1. (Always cot function lies between 0 and 1)

**D2.  $A = B$  if and only if  $IFMS(A, B) = 0$** 

(i) Let the two *IFMS* A and B be equal (i.e.)  $A = B$ . This implies for any  $\mu_A^j(x_i) = \mu_B^j(x_i)$  and  $\vartheta_A^j(x_i) = \vartheta_B^j(x_i)$  which states that  $|\mu_A^j(x_i) - \mu_B^j(x_i)|$  and  $|\vartheta_A^j(x_i) - \vartheta_B^j(x_i)| = 0$ . Hence  $IFMS(A, B) = 0$

(ii) Let the  $IFMS(A, B) = 0$

The zero distance measure is possible only if both  $|\mu_A^j(x_i) - \mu_B^j(x_i)|$  and  $|\vartheta_A^j(x_i) - \vartheta_B^j(x_i)| = 0$ , as the Cotangent similarity measure concerns with max operator ( $\vee$ ) of membership and non membership difference. This refers that  $\mu_A^j(x_i) = \mu_B^j(x_i)$  and  $\vartheta_A^j(x_i) = \vartheta_B^j(x_i)$  for all i, j values. Hence  $A = B$ .

**D3.  $IFMS(A, B) = IFMS(B, A)$** 

It is obvious that  $\mu_A^j(x_i) - \mu_B^j(x_i) \neq \mu_B^j(x_i) - \mu_A^j(x_i)$  and  $\vartheta_A^j(x_i) - \vartheta_B^j(x_i) \neq \vartheta_B^j(x_i) - \vartheta_A^j(x_i)$

But  $|\mu_A^j(x_i) - \mu_B^j(x_i)| = |\mu_B^j(x_i) - \mu_A^j(x_i)|$  and  $|\vartheta_A^j(x_i) - \vartheta_B^j(x_i)| = |\vartheta_B^j(x_i) - \vartheta_A^j(x_i)|$

$$\begin{aligned} \text{Hence } IFMS(A, B) &= \frac{1}{\eta} \sum_{j=1}^{\eta} \left[ \frac{1}{n} \sum_{i=1}^n \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A(x_i) - \mu_B(x_i)| \vee |\vartheta_A(x_i) - \vartheta_B(x_i)|)}{4} \right\} \right] \\ &= \frac{1}{\eta} \sum_{j=1}^{\eta} \left[ \frac{1}{n} \sum_{i=1}^n \text{Cot} \left\{ \frac{\pi + \pi(|\mu_B(x_i) - \mu_A(x_i)| \vee |\vartheta_B(x_i) - \vartheta_A(x_i)|)}{4} \right\} \right] = IFMS(B, A) \end{aligned}$$

**D4. If  $A \subseteq B \subseteq C$  for A, B, C are *IFMS* then  $IFMS(A, B) \leq IFMS(A, C)$  and  $IFMS(B, C) \leq IFMS(A, C)$** 

Let  $A \subseteq B \subseteq C$ , then the assumption is  $\mu_A^j(x_i) \leq \mu_B^j(x_i) \leq \mu_C^j(x_i)$  and  $\vartheta_A^j(x_i) \geq \vartheta_B^j(x_i) \geq \vartheta_C^j(x_i)$  for every  $x_i \in X$

**Case (i)** Let  $|\mu_A^j(x_i) - \mu_C^j(x_i)| \geq |\vartheta_A^j(x_i) - \vartheta_C^j(x_i)|$  Then from the assumption of non membership function, we have  $|\vartheta_A^j(x_i) - \vartheta_B^j(x_i)| \leq |\vartheta_A^j(x_i) - \vartheta_C^j(x_i)| \leq |\mu_A^j(x_i) - \mu_C^j(x_i)|$  ---- (3.4.1)

$$\text{Also } |\vartheta_B^j(x_i) - \vartheta_C^j(x_i)| \leq |\vartheta_A^j(x_i) - \vartheta_C^j(x_i)| \leq |\mu_A^j(x_i) - \mu_C^j(x_i)| \quad \text{---- (3.4.2)}$$

Now from the assumption of the membership, we have

$$|\mu_A^j(x_i) - \mu_B^j(x_i)| \leq |\mu_A^j(x_i) - \mu_C^j(x_i)| \text{ and } |\mu_B^j(x_i) - \mu_C^j(x_i)| \leq |\mu_A^j(x_i) - \mu_C^j(x_i)| \quad \text{---- (3.4.3)}$$

$$\text{From (3.4.1, 3.4.2, 3.4.3) } IFMS(A, B) \leq IFMS(A, C) \text{ and } IFMS(B, C) \leq IFMS(A, C)$$

**Case (ii)** Let  $|\mu_A^j(x_i) - \mu_C^j(x_i)| \leq |\vartheta_A^j(x_i) - \vartheta_C^j(x_i)|$  Then from the assumption of membership function, we

$$\text{have } |\mu_A^j(x_i) - \mu_B^j(x_i)| \leq |\mu_A^j(x_i) - \mu_C^j(x_i)| \leq |\vartheta_A^j(x_i) - \vartheta_C^j(x_i)| \quad \text{---- (3.4.4)}$$

$$\text{Also } |\mu_B^j(x_i) - \mu_C^j(x_i)| \leq |\mu_A^j(x_i) - \mu_C^j(x_i)| \leq |\vartheta_A^j(x_i) - \vartheta_C^j(x_i)| \quad \text{---- (3.4.5)}$$

Now from the assumption of the non membership, we have

$$|\vartheta_A^j(x_i) - \vartheta_B^j(x_i)| \leq |\vartheta_A^j(x_i) - \vartheta_C^j(x_i)| \text{ and } |\vartheta_B^j(x_i) - \vartheta_C^j(x_i)| \leq |\vartheta_A^j(x_i) - \vartheta_C^j(x_i)| \quad \text{-- (3.4.6)}$$

$$\text{From (3.4.4, 3.4.5, 3.4.6) } IFMS(A, B) \leq IFMS(A, C) \text{ and } IFMS(B, C) \leq IFMS(A, C)$$

### NUMERICAL EVALUATION: 3.3

#### EXAMPLE: 4.1

Let  $X = \{A_1, A_2, A_3, A_4, \dots, A_n\}$  with  $A = \{A_1, A_2, A_3, A_4, A_5\}$  and  $B = \{A_6, A_7, A_8, A_9, A_{10}\}$  are the IFMS defined as

$$A = \{ \langle A_1 : (0.6, 0.4), (0.5, 0.5) \rangle, \langle A_2 : (0.5, 0.3), (0.4, 0.5) \rangle, \langle A_3 : (0.5, 0.2), (0.4, 0.4) \rangle, \\ \langle A_4 : (0.3, 0.2), (0.3, 0.2) \rangle, \langle A_5 : (0.2, 0.1), (0.2, 0.2) \rangle \} \\ B = \{ \langle A_6 : (0.8, 0.1), (0.4, 0.6) \rangle, \langle A_7 : (0.7, 0.3), (0.4, 0.2) \rangle, \langle A_8 : (0.4, 0.5), (0.3, 0.3) \rangle, \\ \langle A_9 : (0.2, 0.7), (0.1, 0.8) \rangle, \langle A_{10} : (0.2, 0.6), (0, 0.6) \rangle \}$$

Here, the cardinality  $\eta = 5$  as  $|Mc(A)| = |NMc(A)| = 5$  and  $|Mc(B)| = |NMc(B)| = 5$  and the cotangent similarity measure is

$$\frac{1}{5} \sum_{j=1}^5 \left[ \frac{1}{2} \sum_{i=1}^2 \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A(x_i) - \mu_B(x_i)| \vee |\vartheta_A(x_i) - \vartheta_B(x_i)|)}{4} \right\} \right] = \mathbf{0.5936}$$

#### EXAMPLE: 4.2

Let  $X = \{A_1, A_2, A_3, A_4, \dots, A_n\}$  with  $A = \{A_1, A_2\}$  and  $B = \{A_9, A_{10}\}$  are the IFMS defined as

$$A = \{ \langle A_1 : (0.1, 0.2) \rangle, \langle A_2 : (0.3, 0.3) \rangle \}, B = \{ \langle A_9 : (0.1, 0.2) \rangle, \langle A_{10} : (0.2, 0.3) \rangle \} \\ \text{Here, the cardinality } \eta = 2 \text{ as } |Mc(A)| = |NMc(A)| = 2 \text{ and } |Mc(B)| = |NMc(B)| = 2 \text{ and the cotangent similarity measure is}$$

$$\frac{1}{2} \sum_{j=1}^2 \left[ \frac{1}{1} \sum_{i=1}^1 \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A(x_i) - \mu_B(x_i)| \vee |\vartheta_A(x_i) - \vartheta_B(x_i)|)}{4} \right\} \right] = \mathbf{0.9271}$$

#### EXAMPLE: 4.3

Let  $X = \{A_1, A_2, A_3, A_4, \dots, A_n\}$  with  $A = \{A_1, A_2\}$  and  $B = \{A_3, A_4\}$  are the IFMS defined as

$$A = \{ \langle A_1 : (0.4, 0.2, 0.1), (0.3, 0.1, 0.2) \rangle, \langle A_2 : (0.2, 0.1, 0.2), (0.1, 0.4, 0.3) \rangle, \\ \langle A_2 : (0.6, 0.3, 0), (0.4, 0.5, 0.1) \rangle, \langle A_2 : (0.4, 0.3, 0.2), (0.2, 0.6, 0.2) \rangle \} \\ B = \{ \langle A_3 : (0.5, 0.2, 0.3), (0.4, 0.2, 0.3) \rangle, \langle A_4 : (0.4, 0.1, 0.2), (0.1, 0.1, 0.6) \rangle, \\ \langle A_4 : (0.4, 0.6, 0.2), (0.4, 0.5, 0) \rangle, \langle A_4 : (0.3, 0.4, 0.2), (0.2, 0.4, 0.1) \rangle \}$$

The cardinality  $\eta = 2$  as  $|Mc(A)| = |NMc(A)| = |Hc(A)| = 2$  and  $|Mc(B)| = |NMc(B)| = |Hc(B)| = 2$

Hence, the cotangent similarity measure is

$$\frac{1}{2} \sum_{j=1}^2 \left[ \frac{1}{4} \sum_{i=1}^4 \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A(x_i) - \mu_B(x_i)| \vee |\vartheta_A(x_i) - \vartheta_B(x_i)| \vee |\pi_A(x_i) - \pi_B(x_i)|)}{4} \right\} \right] = \mathbf{0.7450}$$

#### EXAMPLE: 4.4

Let  $X = \{A_1, A_2, A_3, A_4, \dots, A_n\}$  with  $A = \{A_1, A_2\}$  and  $B = \{A_6\}$  such that the IFMS A and B are

$$A = \{ \langle A_1 : (0.6, 0.2, 0.2), (0.4, 0.3, 0.3) \rangle, \langle A_1 : (0.1, 0.7, 0.2) \rangle, \\ \langle A_2 : (0.7, 0.1, 0.2), (0.3, 0.6, 0.1) \rangle, \langle A_2 : (0.2, 0.7, 0.1) \rangle \} \\ B = \{ \langle A_6 : (0.8, 0.1, 0.1), (0.2, 0.7, 0.1) \rangle, \langle A_6 : (0.3, 0.5, 0.2) \rangle \}$$

As  $|Mc(A)| = |NMc(A)| = |Hc(A)| = 2$  and  $|Mc(B)| = |NMc(B)| = |Hc(B)| = 1$ ,

their cardinality  $\eta = \text{Max} \{ \eta(A), \eta(B) \} = \max \{ 2, 1 \} = 2$ . Therefore the cotangent similarity measure is

$$\frac{1}{2} \sum_{j=1}^2 \left[ \frac{1}{3} \sum_{i=1}^3 \text{Cot} \left\{ \frac{\pi + \pi(|\mu_A(x_i) - \mu_B(x_i)| \vee |\vartheta_A(x_i) - \vartheta_B(x_i)| \vee |\pi_A(x_i) - \pi_B(x_i)|)}{4} \right\} \right] = \mathbf{0.7329}$$

## IV MEDICAL DIAGNOSIS USING IFMS - COTANGENT MEASURE

As Medical diagnosis contains lots of uncertainties, they are the most interesting and fruitful areas of application for fuzzy set theory. In some situations, terms of membership function alone is not adequate. Hence, the Intuitionistic fuzzy set theory consisting of both the terms like membership and non membership function is considered to be the better one. Due to the increased volume of information available to physicians from new medical technologies, the process of classifying different set of symptoms under a single name of disease becomes difficult. Recently, there are various models of medical diagnosis under the general framework of fuzzy sets are proposed. In some practical situations, there is the possibility of each element having different membership and non membership functions. The proposed distance and similarity measure among the Patients Vs Symptoms and Symptoms Vs diseases gives the proper medical diagnosis. The unique feature of this proposed method is that it considers multi membership and non membership. By taking one

time inspection, there may be error in diagnosis. Hence, this multi time inspection, by taking the samples of the same patient at different times gives best diagnosis.

Let  $P = \{ P_1, P_2, P_3, P_4 \}$  be a set of Patients.

$D = \{ \text{Fever, Tuberculosis, Typhoid, Throat disease} \}$  be the set of diseases

and  $S = \{ \text{Temperature, Cough, Throat pain, Headache, Body pain} \}$  be the set of symptoms.

Our solution is to examine the patient at different time intervals (three times a day), which in turn give arise to different membership and non membership function for each patient.

**TABLE : 4.1 – IFMs Q : The Relation between Patient and Symptoms**

Q	Temperature	Cough	Throat Pain	Head Ache	Body Pain
P <sub>1</sub>	(0.6, 0.2)	(0.4, 0.3)	(0.1, 0.7)	(0.5, 0.4)	(0.2, 0.6)
	(0.7, 0.1)	(0.3, 0.6)	(0.2, 0.7)	(0.6, 0.3)	(0.3, 0.4)
	(0.5, 0.4)	(0.4, 0.4)	(0, 0.8)	(0.7, 0.2)	(0.4, 0.4)
P <sub>2</sub>	(0.4, 0.5)	(0.7, 0.2)	(0.6, 0.3)	(0.3, 0.7)	(0.8, 0.1)
	(0.3, 0.4)	(0.6, 0.2)	(0.5, 0.3)	(0.6, 0.3)	(0.7, 0.2)
	(0.5, 0.4)	(0.8, 0.1)	(0.4, 0.4)	(0.2, 0.7)	(0.5, 0.3)
P <sub>3</sub>	(0.1, 0.7)	(0.3, 0.6)	(0.8, 0)	(0.3, 0.6)	(0.4, 0.4)
	(0.2, 0.6)	(0.2, 0)	(0.7, 0.1)	(0.2, 0.7)	(0.3, 0.7)
	(0.1, 0.9)	(0.1, 0.7)	(0.8, 0.1)	(0.2, 0.6)	(0.2, 0.7)

Let the samples be taken at three different timings in a day (morning, noon and night)

**TABLE : 4.2 – IFMs R : The Relation among Symptoms and Diseases**

R	Viral Fever	Tuberculosis	Typhoid	Throat disease
Temperature	(0.8, 0.1)	(0.2, 0.7)	(0.5, 0.3)	(0.1, 0.7)
Cough	(0.2, 0.7)	(0.9, 0)	(0.3, 0.5)	(0.3, 0.6)
Throat Pain	(0.3, 0.5)	(0.7, 0.2)	(0.2, 0.7)	(0.8, 0.1)
Head ache	(0.5, 0.3)	(0.6, 0.3)	(0.2, 0.6)	(0.1, 0.8)
Body ache	(0.5, 0.4)	(0.7, 0.2)	(0.4, 0.4)	(0.1, 0.8)

**TABLE : 4.3 – The Cotangent Similarity Measure between IFMs Q and R :**

Cotangent Similarity measure	Viral Fever	Tuberculosis	Typhoid	Throat disease
P <sub>1</sub>	0.7242	0.5152	<b>0.7711</b>	0.4868
P <sub>2</sub>	0.5967	<b>0.7292</b>	0.6446	0.5327
P <sub>3</sub>	0.5271	0.5777	0.6177	<b>0.7825</b>

The *highest similarity measure* from the table 4.3 gives the proper medical diagnosis.

Patient **P1** suffers from **Typhoid**, Patient **P2** suffers from **Tuberculosis** and Patient **P3** suffers from **Throat disease**.

## V. CONCLUSION

In this paper, we have developed the method for deriving the cotangent similarity measure of *IFMS* from *IFS* theory. The prominent characteristic of this method is that it considers multi membership, non membership functions and this similarity measure also guarantee that the cotangent function of any two *IFMSs* equals to

one if and only if the two *IFMSs* are the same. The **example 3.1, 3.2** shows that the new measure perform well in the case of membership and non membership function and **example 3.3, 3.4** depicts that the proposed measure is effective with three representatives of *IFMSs* – membership, non membership and hesitation functions. Hence, from the analysis and numerical evaluation, it is clear that this proposed method can be applied to any pattern recognitions and decision making problems. Finally, an illustrative example, the medical diagnosis has been given to show the efficiency of the developed cotangent similarity measure.

## REFERENCES

- [1] **Atanassov K.**, Intuitionistic fuzzy sets, *Fuzzy Sets and System* 20 (1986) 87-96.
- [2] **Atanassov K.**, More on Intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 33 (1989) 37-46.
- [3] **Zadeh L. A.**, *Fuzzy sets, Information and Control* 8 (1965) 338-353.
- [4] **Blizard W. D.**, Multi set Theory, *Notre Dame Journal of Formal Logic, Vol. 30, No. 1* 36-66, (1989).
- [5] **Yager R. R.**, On the theory of bags,(Multi sets), *Int. Jou. Of General System*, 13 (1986) 23-37.
- [6] **Shinoj T.K., Sunil Jacob John** , Intuitionistic Fuzzy Multi sets and its Application in Medical Diagnosis, *World Academy of Science, Engineering and Technology, Vol. 61* (2012).
- [7] **Chen S.M., Tan J.M.**, Handling multi-criteria fuzzy decision-making problems based on vague sets. *Fuzzy sets and systems*, 67 (1994), 163-172.
- [8] **Szmidt E., Kacprzyk J.**, On measuring distances between Intuitionistic fuzzy sets, *Notes on IFS, Vol.3*(1997) 1- 13.
- [9] **Szmidt E., Kacprzyk J.**, Distances between Intuitionistic fuzzy sets. *Fuzzy Sets System*, 114 (2000) 505-518.
- [10] **Szmidt E., Kacprzyk J.**, Entropy for Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, 118 (2001) 467-477.
- [11] **Szmidt E., Kacprzyk J.**, A similarity measure for Intuitionistic fuzzy sets and its application in supporting medical diagnostic reasoning. *ICAISC 2004, Vol. LNAI 3070* (2004) 388-393.
- [12] **Xu Z. S., Xia M.M.**, Some new similarity measures for Intuitionistic fuzzy values and their application in group decision making *Journal of System Science Engin*, 19 (2010) 430-452.
- [13] **Wang J. Q., Wang P.**, Intuitionistic linguistic fuzzy multi-criteria decision making based on Intuitionistic fuzzy entropy. *Control and Decision*, 27(11) (2012) 1694-1968.
- [14] **Tian Maoying .,** A new fuzzy similarity based on cotangent Function for medical diagnosis, *Advanced Modeling and Optimization* , Vol. 15 Number 2 (2013).
- [15] **Rajarajeswari P., Uma N.**, On Distance and Similarity Measures of Intuitionistic Fuzzy Multi Set, *IOSR Journal of Mathematics (IOSR-JM)*, Vol. 5, Issue 4 (Jan. - Feb. 2013) PP 19-23.
- [16] **Rajarajeswari P., Uma N.**, Hausdroff Similarity measures for Intuitionistic Fuzzy Multi Sets and Its Application in Medical diagnosis *International Journal of Mathematical Archive-4*(9) (2013) 106-111.